arXiv:math/0405423v1 [math.NT] 21 May 2004

A conjecture-generalization of Sondow's formula

Petros Hadjicostas
Department of Mathematics and Statistics
Texas Tech University, Box 41042
Lubbock, TX 79409-1042, USA

By generalizing Beukers' arguments in [2] one can easily deduce that (e.g., see [6]):

Theorem 1 *Let* $\mathbb{N}$ *be the set of all non-negative integers, and for all* $n \in \mathbb{N}\backslash\{0\}$ *let* d_n *be the lowest common multiple of* $1, 2, \ldots, n$. *Let* $r, s \in \mathbb{N}$.

(a) If $r > s$, *then for all* $n \in \mathbb{N}$,

$$\int_0^1 \int_0^1 \frac{\ln^n(xy)}{1-xy} x^r y^s \, dx\, dy = \frac{n!(-1)^n}{r-s} \sum_{k=s+1}^{r} \frac{1}{k^{n+1}},$$

which is a rational number whose denominator is a divisor of d_r^{n+2}.

(b) If $r = s$, *then for all* $n \in \mathbb{N}$,

$$\int_0^1 \int_0^1 \frac{\ln^n(xy)}{1-xy} x^r y^r \, dx\, dy = (n+1)!(-1)^n \left(\zeta(n+2) - \sum_{k=1}^{r} \frac{1}{k^{n+2}} \right).$$

(Note that the sum of numbers over the empty set is by definition zero.)

(For related results see also the papers [1], [5], [7], [8], [9], [10], [12], [15], and [16]. See also the books [3, pp. 365-370] and [4, Appendix A2, pp. 372-381].) Using parts (a) and (b) of Theorem 1, it is easy to show that

$$\int_0^1 \int_0^1 \frac{[-\ln(xy)]^n}{1-xy} (1-x)\, dx\, dy = \Gamma(n+2) \left[\zeta(n+2) - \frac{1}{n+1} \right],$$

for all $n \in \mathbb{N}$. This gives rise to the following conjecture:

Conjecture 1 *For* $z \in \mathbb{C}$ *with* $\Re(z) > -2$,

$$\int_0^1 \int_0^1 \frac{[-\ln(xy)]^z}{1-xy} (1-x)\, dx\, dy = \Gamma(z+2) \left[\zeta(z+2) - \frac{1}{z+1} \right]. \tag{0.1}$$

The conjecture can *probably* be proven by generalizing Beukers' arguments in [2] and by using Fractional Calculus (e.g., see [11]). By taking the limit as $z \to -1$ on both sides of (0.1), one obtains Sondow's formula

$$\gamma = \int_0^1 \int_0^1 -\frac{1-x}{(1-xy)\ln(xy)}\, dx\, dy,$$

where γ is Euler's constant. (See [13] and [14].)

References

[1] Ball, K., and Rivoal, T. (2001), "Irrationalité d'une infinité de valeurs de la fonction zêta aux entiers impairs (The irrationality of an infinite number of values of the zeta function at odd integers)," *Invent. Math.*, **146**, 193–207.

[2] Beukers, F. (1979), "A note on the irrationality of $\zeta(2)$ and $\zeta(3)$," *Bull. London Math. Soc.*, **11**, 268-272.

[3] Borwein, J.M., and Borwein, P.B., *Pi and the AGM*, John Wiley, New York, 1987.

[4] Borwein, P., and Erdélyi, T., *Polynomials and polynomial inequalities*, Springer, New York, 1995.

[5] Dvornicich, R., and Viola, C. (1990), "Some remarks on Beukers' integrals," *Number theory*, Vol. 2 (Budapest, 1987), *Colloq. Math. Soc. János Bolyai*, **51**, North-Holland, Amsterdam, 637-657.

[6] Hadjicostas, P. (2002), "Some generalizations of Beukers' integrals," *Kyungpook Mathematical Journal*, **42**, 399-416.

[7] Hata, M. (1995), "A note on Beukers' integral," *J. Austral. Math. Soc. Ser. A*, **58**, 143–153.

[8] Huylebrouck, D. (2001), "Similarities in irrationality proofs for π, $\ln 2$, $\zeta(2)$, and $\zeta(3)$," *Amer. Math. Monthly*, **108**, 222-231.

[9] Krattenthaler, C., and Rivoal, T. (2003), "An identity of Andrews, multiple integrals, and very-well-poised hypergeometric series," *(http://arXiv.org/abs/math/0312148)*, submitted for publication.

[10] Nesterenko, Yu. V. (2003), "Integral identities and constructions of approximations to zeta-values," Actes des 12èmes rencontres arithmétiques de Caen (June 29–30, 2001), *J. Théorie Nombres Bordeaux*, **15**, 535–550.

[11] Oldham, K.B., and Spanier, J., *The Fractional Calculus*, Academic Press, New York, 1974.

[12] Rivoal, T. (2000), "La fonction zêta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs (There are infinitely many irrational values of the Riemann zeta function at odd integers)," *C. R. Acad. Sci. Paris Sér. I Math.*, **331**, 267-270.

[13] Sondow, J. (2004), "A faster product for π and a new integral for $\ln(\pi/2)$," to appear in *The American Mathematical Monthly, (http://arXiv.org/abs/math.NT/0401406).*

[14] Sondow, J. (2004), "Double integrals for Euler's constant and $\ln(4/\pi)$," submitted to the *The American Mathematical Monthly, (http://arXiv.org/abs/math.CA/0211148).*

[15] Zudilin, W. (2002), "Very well-poised hypergeometric series and multiple integrals," *Russian Math. Surveys*, **57**, 824–826.

[16] Zudilin, W. (2004), "Well-poised hypergeometric transformations of Euler-type multiple integrals," *J. London Math. Soc. (2)*, **70**, 16 pages.